

Fermionic Warm Dark Matter reproduces galaxy observations because of quantum mechanics

H. J. de Vega

LPTHE, CNRS/Université P & M Curie (Paris VI).

**Workshop CIAS Meudon 2013: WARM DARK MATTER GALAXIES IN AGREEMENT
WITH OBSERVATIONS**

Observatoire de Paris, Château de Meudon, Meudon campus 5, 6 and 7 June 2013

Dark Matter in the Universe

81 % of the matter of the universe is **DARK** (DM).
DM is the dominant component of galaxies.

DM interacts through **gravity**.

Further DM interactions **unobserved** so far. Such couplings must be **very weak**: much weaker than weak interactions.

DM is **outside** the standard model of particle physics.

Proposed candidates:

- Neutrinos: HDM, (in the 1980's) $m \sim 1$ eV. Ruled out !
- Cold Dark Matter: CDM, WIMPS, $m \sim 10 - 1000$ GeV.
IN BIG TROUBLE.
- Warm Dark Matter: WDM, sterile neutrinos $m \sim 1$ keV.
THE ANSWER !

Dark Matter Particles

DM particles decouple due to the universe expansion, their distribution function **freezes out** at decoupling.

Early decoupling: $T_d \sim 100 \text{ GeV}$

The characteristic length scale is the **free streaming scale** (or Jeans' scale). For DM particles decoupling UR:

$$r_{Jeans} = 57.2 \text{ kpc} \frac{\text{keV}}{m} \left(\frac{100}{g_d} \right)^{\frac{1}{3}}, \text{ solving the linear Boltz-V eqs.}$$

g_d = number of UR degrees of freedom at decoupling.

DM particles can **freely** propagate over distances of the order of the free streaming scale.

Therefore, structures at scales smaller or of the order of r_{Jeans} are **erased**.

The size of the DM galaxy cores is in the tenths of kpc $\Rightarrow m$ should be in the keV scale (Warm Dark Matter particles, WDM).

CDM free streaming scale

For CDM particles with $m \sim 100 \text{ GeV} \Rightarrow r_{\text{Jeans}} \sim 0.1 \text{ pc}$.

Hence CDM structures keep forming till scales as small as the solar system.

This is a **robust result** of N -body CDM simulations but **never observed** in the sky. Including baryons do not cure this serious problem. There is **over abundance** of small structures in CDM (also called the satellite problem).

CDM has **many serious** conflicts with observations:

Galaxies naturally grow through merging in CDM models.

Observations show that galaxy mergers are **rare** ($< 10\%$).

Pure-disk galaxies (bulgeless) are observed whose formation through CDM is **unexplained**.

CDM predicts **cusped** density profiles: $\rho(r) \sim 1/r$ for small r .

Observations show **cored** profiles: $\rho(r)$ bounded for small r .

Adding by hand **strong** enough feedback from baryons can eliminate cusps but **spoils** the star formation rate.

Structure Formation in the Universe

Structures in the Universe as galaxies and cluster of galaxies form out of the **small primordial quantum fluctuations** originated by inflation just after the big-bang.

These linear small primordial fluctuations **grow** due to gravitational unstabilities (Jeans) and then classicalize.

Structures form through non-linear gravitational evolution.

Hierarchical formation starts from small scales first.


N -body CDM simulations **fail** to produce the observed structures for **small** scales less than some kpc.

Both N -body WDM and CDM simulations yield **identical and correct** structures for scales larger than some kpc.

WDM predicts **correct structures for small scales** (below kpc) when its **quantum** nature is taken into account.

Primordial power $P(k)$: first ingredient in galaxy formation.

Linear primordial power today $P(k)$ vs. k Mpc h



$\log_{10} P(k)$ vs. $\log_{10}[k \text{ Mpc } h]$ for **CDM**, **1 keV**, **2 keV**,
light-blue 4 keV DM particles decoupling in equil, and 1
keV **sterile neutrinos**. WDM cuts $P(k)$ on small scales
 $r \lesssim 100 (\text{keV}/m)^{4/3} \text{ kpc}$. CDM and WDM identical for CMB.

Summary Warm Dark Matter, WDM: $m \sim \text{keV}$

- Large Scales, structures beyond ~ 100 kpc: WDM and CDM yield **identical** results **which agree with observations**
 - Intermediate Scales: WDM give the **correct abundance** of substructures.
 - Inside galaxy cores, below ~ 100 pc: N-body classical physics simulations are **incorrect** for WDM because of **important quantum effects**.
 - Quantum calculations (Thomas-Fermi) give galaxy cores, galaxy masses, velocity dispersions and densities in **agreement with the observations**.
 - Direct Detection of the main WDM candidate: the sterile neutrino. **Beta decay and electron capture**. ^3H , Re, Ho.
- So far, **not a single valid** objection arose against WDM.
- Baryons (=16%DM) expected to give a correction to WDM

Quantum physics in Galaxies

de Broglie wavelength of DM particles $\lambda_{dB} = \frac{\hbar}{m v}$

d = mean distance between particles, v = mean velocity

$$d = \left(\frac{m}{\rho}\right)^{\frac{1}{3}}, \quad Q = \rho/v^3, \quad Q = \text{phase space density.}$$

ratio: $\mathcal{R} = \frac{\lambda_{dB}}{d} = \hbar \left(\frac{Q}{m^4}\right)^{\frac{1}{3}}$

Observed values: $2 \times 10^{-3} \left(\frac{\text{keV}}{m}\right)^{\frac{4}{3}} < \mathcal{R} < 1.4 \left(\frac{\text{keV}}{m}\right)^{\frac{4}{3}}$

The **larger** \mathcal{R} is for ultracompact dwarfs.

The **smaller** \mathcal{R} is for big spirals.

\mathcal{R} near unity (or above) means a **QUANTUM OBJECT**.

Observations alone show that compact dwarf galaxies are **quantum objects** (for WDM).

No quantum effects in CDM: $m \gtrsim \text{GeV} \Rightarrow \mathcal{R} \lesssim 10^{-8}$

Quantum pressure vs. gravitational pressure

quantum pressure: $P_q = \text{flux of momentum} = n v p$,

$v = \text{mean velocity}$, momentum $= p \sim \hbar / \Delta x \sim \hbar n^{\frac{1}{3}}$,

particle number density $= n = \frac{M_q}{\frac{4}{3} \pi R_q^3 m}$

galaxy mass $= M_q$, galaxy halo radius $= R_q$

gravitational pressure: $P_G = \frac{G M_q^2}{R_q^2} \times \frac{1}{4 \pi R_q^2}$

Equilibrium: $P_q = P_G \implies$

$$R_q = \frac{3^{\frac{5}{3}}}{(4 \pi)^{\frac{2}{3}}} \frac{\hbar^2}{G m^{\frac{8}{3}} M_q^{\frac{1}{3}}} = 10.6 \dots \text{pc} \left(\frac{10^6 M_{\odot}}{M_q} \right)^{\frac{1}{3}} \left(\frac{\text{keV}}{m} \right)^{\frac{8}{3}}$$

$$v = \left(\frac{4 \pi}{81} \right)^{\frac{1}{3}} \frac{G}{\hbar} m^{\frac{4}{3}} M_q^{\frac{2}{3}} = 11.6 \frac{\text{km}}{\text{s}} \left(\frac{\text{keV}}{m} \right)^{\frac{4}{3}} \left(\frac{M_q}{10^6 M_{\odot}} \right)^{\frac{2}{3}}$$

for WDM the values of M_q , R_q and v are **consistent with the dwarf galaxy observations !!**.

Dwarf spheroidal galaxies **can be supported** by the fermionic quantum pressure of WDM.

Self-gravitating Fermions in the Thomas-Fermi approach

WDM is non-relativistic in the MD era. A single DM halo in late stages of formation relaxes to a time-independent form especially in the interior.

Chemical potential: $\mu(r) = \mu_0 - m \phi(r)$, $\phi(r) = \text{grav. pot.}$

Poisson's equation: $\frac{d^2\mu}{dr^2} + \frac{2}{r} \frac{d\mu}{dr} = -4\pi G m \rho(r)$

$\rho(0) = \text{finite for fermions} \implies \frac{d\mu}{dr}(0) = 0.$

Density $\rho(r)$ and pressure $P(r)$ in terms of the distribution function $f(E)$:


$$\rho(r) = \frac{m}{\pi^2 \hbar^3} \int_0^\infty p^2 dp f\left[\frac{p^2}{2m} - \mu(r)\right]$$

$$P(r) = \frac{m}{3\pi^2 \hbar^3} \int_0^\infty p^4 dp f\left[\frac{p^2}{2m} - \mu(r)\right]$$

Boundary condition at

$$r = R = R_{200} \sim R_{vir} , \quad \rho(R_{200}) \simeq 200 \bar{\rho}_{DM}$$


Q vs. halo radius. Galaxy observations vs. Thomas-Fermi



observed $Q = \rho/v^3$ from stars are **upper bounds** for DM Q

Density and velocity profiles from Thomas-Fermi

Cored density profile and velocity profile obtained from Thomas-Fermi.



Galaxy data vs. Thomas-Fermi

Mass, halo radius, velocity dispersion and central density from a broad variety of galaxies: ultracompact galaxies to giant spirals, Willman 1, Segue 1, Canis Venatici II, Coma-Berenices, Leo II, Leo T, Hercules, Carina, Ursa Major I, Draco, Leo I, Sculptor, Boötes, Canis Venatici I, Sextans, Ursa Minor, Fornax, NGC 185, NGC 855, NGC 4478, NGC 731, NGC 3853, NGC 499 and a large number of spiral galaxies.

Phase-Space distribution function $f(E/E_0)$: Fermi-Dirac ($F(x) = \frac{1}{e^x + 1}$) and out of equilibrium sterile neutrinos give similar results.

E_0 = effective galaxy temperature (energy scale).

E_0 turns to be $10^{-3} \text{ } ^\circ\text{K} < E_0 < 50 \text{ } ^\circ\text{K}$

colder = ultracompact, warmer = large spirals.

$E_0 \sim m < v^2 >_{\text{observed}}$ for $m \sim 2 \text{ keV}$.

Self-gravitating Fermions in the Thomas-Fermi approach

The Thomas-Fermi approach gives physical galaxy magnitudes: mass, halo radius, phase-space density and velocity dispersion **fully compatible** with observations from the largest spiral galaxies till the ultracompact dwarf galaxies for a WDM particle mass **around 2 keV**.


Compact dwarf galaxies are close to a degenerate WDM Fermi gas while large galaxies are classical WDM Boltzmann gases.

Thomas-Fermi approach **works in the classical (Boltzmann) regime** too: we always obtain cores with observed sizes.

Fermionic WDM **treated quantum mechanically is able to reproduce** the observed galaxies.

C. Destri, H. J. de Vega, N. G. Sanchez,
arXiv:1204.3090, New Astronomy **22**, 39 (2013) and
arXiv:1301.1864, Astroparticle Physics, **46**, 14 (2013).


Velocity widths in galaxies: test substructure formation



Velocity widths in galaxies from 21cm HI surveys. ALFALFA survey **clearly favours WDM** over CDM. (Papastergis et al. ApJ, 2011, Zavala et al. ApJ, 2009).

Notice that the WDM **red** curve is for $m = 1 \text{ keV}$ WDM particle decoupling at thermal equilibrium.

The 1 keV WDM curve falls somehow below the data suggesting a slightly **larger** WDM particle mass.



Small scale structures at high redshift.


WDM (green continuous line) **reproduces** the observed small scale structures for redshifts up to eight where observations are available.

Danese, de Vega, Lapi, Salucci, Sanchez (in preparation).

The evolution of the AGN luminosity function in WDM vs CDM

WDM: shaded grey area CDM red dashed line

NM , Fiore, Lamastra 2012b



Sterile Neutrinos $\nu_s \simeq \nu_R + \theta \nu_L$

Sterile neutrinos ν_s : named by Bruno Pontecorvo (1968).

Singlets under all SM symmetries.

Do not interact weak, neither EM, nor strongly.

WDM ν_s can be produced from active neutrinos by mixing.

Mixing angles: $\theta \sim 10^{-3} - 10^{-4}$ (depending on the model) are appropriate **to produce enough** ν_s accounting for the observed total DM.

Smallness of θ makes sterile neutrinos **difficult** to detect.

Sterile neutrinos **can be detected** in beta decay and in electron capture (EC) when a ν_s with mass in the keV scale is produced **instead** of an active ν_e .

Beta decay: the electron spectrum is slightly modified at energies around the mass (\sim keV) of the ν_s .

$${}^3H_1 \Longrightarrow {}^3He_2 + e^- + \bar{\nu}_e \quad , \quad {}^{187}Re \Longrightarrow {}^{187}Os + e^- + \bar{\nu}_e.$$

The electron energy spectrum is observed.

Electron Capture and Sterile Neutrinos

Electron capture: $^{163}\text{Ho} + e^- \Longrightarrow ^{163}\text{Dy}^* + \nu_e$

The nonradiative de-excitation of the Dy^* is observed and is different for ν_s in the keV range than for active ν_e .

Available energies:

$Q(^{187}\text{Re}) = 2.47 \text{ keV}$, $Q(^3\text{H}_1) = 18.6 \text{ keV}$, $Q(^{163}\text{Ho}) \simeq 2.5 \text{ keV}$.

Theoretical analysis of ν_s detection in Rhenium and Tritium beta decay: H J de V, O. Moreno, E. Moya, M. Ramón Medrano, N. Sánchez, Nucl. Phys. B866, 177 (2013).

Present experiments searching the small active neutrino mass also look for sterile neutrinos in the keV scale:

MARE (Milan, Italy), Rhenium beta decay and Holmiun EC.

KATRIN (Karlsruhe, Germany), Tritium beta decay.

ECHo (Heidelberg, Germany), Holmiun EC.

Project 8, (Seattle, USA) Tritium beta decay (still in project).

Sterile neutrino models

- DW: Dodelson-Widrow model (1994) sterile neutrinos produced by non-resonant mixing from active neutrinos.
- Shi-Fuller model (1998) sterile neutrinos produced by resonant mixing from active neutrinos.
- ν MSM model (2005) sterile neutrinos produced by a Yukawa coupling from a real scalar χ .
- Models based on: Froggatt-Nielsen mechanism, flavor symmetries, Q_6 , split see-saw, extended see-saw, inverse see-saw, loop mass. Furthermore: scotogenic, LR symmetric, etc. Review by A Merle (2013).

WDM particles in the first 3 models behave primordially just as if their masses were different (FD = thermal fermions):

$$\frac{m_{DW}}{\text{keV}} \simeq 2.85 \left(\frac{m_{FD}}{\text{keV}} \right)^{\frac{4}{3}}, \quad m_{SF} \simeq 2.55 m_{FD}, \quad m_{\nu\text{MSM}} \simeq 1.9 m_{FD}.$$

H J de Vega, N Sanchez, Warm Dark Matter cosmological fluctuations, Phys. Rev. D85, 043516 and 043517 (2012).

X-ray detection of DM sterile neutrinos

Sterile neutrinos ν_s decay into active neutrinos ν_e plus **X-rays** with a lifetime $\sim 10^{11} \times$ age of the universe.

These X-rays **may be seen** in the sky looking to galaxies !


recent review: C. R. Watson et al. JCAP, (2012).

Future observations:

- DM bridge between M81 and M82 ~ 50 kpc. Overlap of DM halos.
Satellite projects: Xenia (NASA), ASTRO-H (Japan).
- **CMB**: WDM decay distorts the blackbody CMB spectrum. The projected PIXIE satellite mission (A. Kogut et al.) can measure WDM sterile neutrino mass.

Results from **Supernovae**: θ unconstrained, $1 < m < 10$ keV, (G. Raffelt & S. Zhou, PRD 2011).

Constraints on the sterile neutrino mass and mixing angle



Dashed = Shi-Fuller model. Dotted = Dodelson-Widrow for fermion asymmetry $L = 0.1, 0.01$ and 0.003 .

Allowed sterile neutrino region in the right lower corner.

Main difficulty: to distinguish the sterile neutrino decay X-ray from narrow X-ray lines emitted by hot ions as Fe.

Direct searches of CDM particles

All direct searches of wimps look for $m \gtrsim 1$ GeV.

Past, present and future reports of signals in such wimp experiments **cannot be due to DM detection** because the DM particle mass is in the keV scale.

The inconclusive signals in such experiments should be originated by **other kinds of phenomena**.

Contradictions between supposed detection signals in DAMA, CDMS-II, CoGeNT, CRESST, XENON100.

CONCLUSION: These signals are **unrelated to DM**.

e^+ and \bar{p} excess in cosmic rays reported by Pamela and Fermi may be explained by astrophysics: P. L. Biermann et al. PRL (2009), P. Blasi, P. D. Serpico PRL (2009).

AMS02: precise measure of the positron fraction in Galactic cosmic rays (CR) (PRL 2013) that increases with energy. Blum, Katz & Waxman (arXiv:1305.1324) show that this is consistent with positron production by the collision of high energy primary CR with the interstellar medium.

Sterile neutrinos and CMB fluctuations

CMB data give the **effective** number of neutrinos, N_{eff} .

N_{eff} is related in a **subtle** way to the number of active neutrinos (3) plus the number of sterile neutrinos.

Planck result: $N_{\text{eff}} = 3.5 \pm 0.5$ (95%; P+WP+highL+ H_0 +BAO)

Entropy conservation determines the contributions to N_{eff} .

WDM sterile neutrino contribution at recombination

$$\Delta N^{WDM} = \left(\frac{T_d}{T_{rc}} \right)^4 = \left[\frac{g_{rc}}{g(T_d)} \right]^{4/3}. \text{ At recombination } g_{rc} = 29/4.$$

WDM decouples early at T_d **beyond** the Fermi scale

The number of UR degrees of freedom at decoupling $g(T_d)$ includes **all SM particles** and probably beyond.

$$g_{SM} = 427/4, \quad g_{MSSM} = 915/4,$$

$$\Delta N_{SM}^{WDM} = 0.02771 \dots, \quad \Delta N_{MSSM}^{WDM} = 0.01003 \dots$$

Too small to be measurable at present !

Conclusion: Planck results say nothing about WDM.

Besides, Planck results are **compatible** with one or two eV sterile neutrinos (see e. g. G. Steigman, 1303.0049).

Summary: keV scale DM particles

- **Reproduce** the phase-space density observed in dwarf spheroidal and spiral galaxies (de Vega, Sanchez, MNRAS 2010).
- Fermionic WDM treated **quantum mechanically** reproduces the main physical galaxy magnitudes: mass, core radius, phase-space density, velocity dispersion, fully consistent with observations and points to a DM particle mass ~ 2 keV (Destri, de Vega, Sanchez, New Astronomy 2012, and 2013).
- The galaxy surface density $\mu_0 \equiv \rho_0 r_0$ is **universal** up to $\pm 10\%$ according to the observations. Its value $\mu_0 \simeq (18 \text{ MeV})^3$ is reproduced by WDM (de Vega, Salucci, Sanchez, New Astronomy, 2012). CDM simulations give 1000 times the observed value of μ_0 (Hoffman et al. ApJ 2007).

Summary: keV scale DM particles

- **Alleviate** the CDM **satellite** problem (Avila-Reese et al. 2000, Götz & Sommer-Larsen 2002, Markovic et al. JCAP 2011) and the CDM **voids** problem (Tikhonov et al. MNRAS 2009).
- Velocity widths in galaxies from 21cm HI surveys. ALFALFA survey clearly favours WDM over CDM. Papastergis et al. ApJ 2011, Zavala et al. ApJ 2009
- Highlights and Conclusions of the **Chalonge Meudon Workshop 2011**: Warm dark matter in the galaxies, arXiv:1109.3187, the **16th Paris Cosmology Colloquium 2011**, arXiv:1203.3562, HJdeV, NGS, and the **Chalonge Meudon Workshop 2012**, arXiv:1305.7452, PLB, HJdeV, NGS.
- 18th Paris Cosmology Colloquium 2013, 24-26 July, Observatoire de Paris.

Future Perspectives

WDM particle models must explain the baryon asymmetry of the universe. An appealing **mass** neutrino hierarchy appears:

- Active neutrino: \sim mili eV
- Light sterile neutrino: \sim eV
- Dark Matter: \sim keV
- Unstable sterile neutrino: \sim MeV....

Need WDM simulations showing substructures, galaxy formation and evolution including **quantum** dynamical evolution. **Quantum** pressure must be included !

WDM simulations should be performed matching semiclassical Hartree-Fock (Thomas-Fermi) dynamics in regions where $Q/m^4 > 0.1$ with classical evolution in regions where $Q/m^4 \ll 1$. Not easy but unavoidable!

Future Perspectives: Detection!

Sterile neutrino detection depends **upon** the particle physics model. There are sterile neutrino models where the keV sterile is **stable** and thus hard to detect.

Astronomical observation of steriles:
X-ray data from galaxy halos.

Direct detection of steriles in Lab:

Bounds on mixing angles from
Mare, Katrin, ECHo and Project 8 are expected.

For a **particle detection** a **dedicated** beta decay or electron capture experiment looks **necessary** to search sterile neutrinos with mass around 2 keV.

Calorimetric techniques seem **well suited**.

Best nuclei for study:

Electron capture in ^{163}Ho , beta decay in ^{187}Re and Tritium.

Richard P. Feynman foresaw the necessity to include quantum physics in simulations in 1981

“I’m not happy with all the analyses that go with just the classical theory, because nature isn’t classical, dammit, and if you want to make a simulation of nature, you’d better make it quantum mechanical, and by golly it’s a wonderful problem, because it doesn’t look so easy.”

Feynman again:

“It doesn’t matter how beautiful your theory is, it doesn’t matter how smart you are. If it doesn’t agree with experiment, it’s wrong.


R. P. Feynman”

**THANK YOU VERY MUCH
FOR YOUR ATTENTION!!**

The expected overdensity


The expected overdensity within a radius R in the linear regime

$$\sigma^2(R) = \int_0^\infty \frac{dk}{k} \Delta^2(k) W^2(kR) \quad , \quad W(kR) : \text{window function.}$$



$\log \sigma(R)$ vs. $\log(R/h \text{ Mpc})$ for CDM, 1 keV, 2 keV, 4 keV DM particles decoupling in equil, and 1 keV (light-blue) sterile neutrinos. WDM flattens and reduces $\sigma(R)$ for small scales.

N-body WDM Simulations: substructure formation



Wednesday, 15 June 2011

WDM subhalos are **less concentrated** than CDM subhalos.

WDM subhalos have the **right concentration** to host the bright Milky Way satellites. Lovell et al. MNRAS (2012).

Summary: WDM produces **correct substructure abundance**.