VOYAGE A TRAVERS L'UNIVERS DE SES ORIGINES A NOS JOURS









Ecole Internationale Daniel Chalonge SESSION OUVERTE d'AUTOMNE 2013 LAST NEWS OF THE UNIVERSE ANTICIPATED



Latest News of the Universe Anticipated

Norma G. SANCHEZ
DR CNRS, LERMA Observatoire de Paris



Ecole Internationale Daniel Chalonge

Observatoire de Paris, Paris 28 /11/2013

CONTENT OF THE UNIVERSE

ATOMS, the building blocks of stars and planets: represent only the $\frac{4.6\%}{}$

DARK MATTER comprises 23.4 % of the universe. This matter, different from atoms, does not emit or absorb light. It has only been detected indirectly by its gravity.

72% of the Universe, is composed of <u>DARK ENERGY</u> that acts as a sort of an anti-gravity.

This energy, distinct from dark matter, is responsible for

the present-day acceleration of the universal expansion, compatible with cosmological constant

CONTENTS

(I) The Standard Model of the Universe Includes Inflation

(II) THE NATURE OF DARK MATTER IN GALAXIES from Theory and Observations: Warm (keV scale) DM

(III) NEW: THE ESSENTIAL ROLE OF QUANTUM PHYSICS IN WDM GALAXIES:

Semiclassical framework: Analytical Results and Numerical (including analytical) Results

Observed Galaxy cores and structures from Fermionic WDM and more results.

(IV) NEW: The generic Galaxy types and properties from a same physical framework: From quantum (compact, dwarfs) to classical (dilute, large) galaxies. Equation of state

HIGHLIGHTS

(I) The Effective (Ginsburg-Landau) Theory of Inflation **PREDICTIONS**:

The Primordial Cosmic Banana: non-zero amount of primordial gravitons. And Forecasts for CMB exps.

(II): TURNING POINT IN THE DARK MATTER PROBLEM: DARK MATTER IN GALAXIES from Theory and Observations: Warm (keV scale) dark matter

Physical Clarification and Simplification
GALAXY FORMATION AND EVOLUTION IN
AGREEMENT WITH OBSERVATIONS
naturally re-insert in COSMOLOGY (LWDM)
Analytical Results and Numerical

NEW RESULTS

FERMIONIC QUANTUM WDM and GRAVITATION DETERMINE THE OBSERVED PHYSICAL GALAXY PROPERTIES

- -> Dark matter (DM) is the main component of galaxies. Quantum mechanics is a cornerstone of physics from microscopic to macroscopic systems as quantum liquids He^3, white dwarf stars and neutron stars.
- -> Recent study: Destri, de Vega, Sanchez, (New Astronomy 22, 39, 2013) suggest that quantum mechanics is also responsible of galaxy structures at the kpc scales and below: near the galaxy center, below 10 100 pc, the DM quantum effects are important for warm DM (WDM), that is for DM particles with masses in the keV scale.
 - -> A new approach to galaxy structure with results in remarkable agreement with observations:

- (i) Dwarf galaxies turn to be quantum macroscopic objects for WDM supported against gravity by the WDM fermion pressure
- (ii) Theoretical analytic framework based on Thomas-Fermi approach determine galaxy structure from the most compact dwarf galaxies to the largest dilute galaxies (spirals, ellipticals).
- The obtained galaxy mass, halo radius, phase-space density, velocity dispersion, are fully consistent with observations.
- (iii) Interestingly enough, a minimal galaxy mass and minimal velocity dispersion are found for DM dominated objects, which in turn imply an universal minimal mass m_min = 1.9 keV for the WDM particle.


- OBSERVED GALAXY CORES vs CDM CUSPS and WDM CORES-
- Astronomical observations show that the DM galaxy density profiles are cored, that is, profiles which are flat at the center.

On the contrary, N-body CDM simulations exhibit cusped density profiles, with a typical 1/r cusped behaviour near the galaxy center r=0.

Classical N-body WDM simulations exhibit cores but with sizes much smaller than the observed cores.

We have recently developped a new approach to this problem thanks to **Quantum Mechanics.**

- Fermions always provide a non vanishing pressure of quantum nature due to the combined action of the Pauli exclusion principle and Heisenberg uncertainty principle.
- Quantum effects for WDM fermions <u>rule out</u> the presence of galaxy cusps for WDM and <u>enlarge</u> the classical core sizes because their <u>repulsive and non-local</u> nature extend well beyond the small pc scales.



From WMAP9 to Planck

Understanding the direction in which data are pointing:

• PREDICTIONS for Planck

- Standard Model of the Universe
 - Standard Single field Inflation
- NO RUNNING of the Primordial Spectral Index
 - NO Primordial NON GAUSSIANITY
 - Neff neutrinos : --> Besides meV active neutrinos:
 - 1 or 2 sterile neutrinos
 - Would opens the sterile neutrino Family:
 - keV sterile neutrino –WDM-

Large Hadron Collider

• The first LHC results at 7-8 TeV, with the discovery of a candidate Higgs boson and the non observation of new particles or exotic phenomena, have made a big step towards completing the experimental confirmation of the Standard Model of particle physics.

• It is thus a good moment to recall our scientific predictions made several years ago on this matter because they are of full actuality.

Large Hadron Collider - LHC-

The results are completely in line with the Standard Model.

No evidence of SUSY at LHC

"Supersymmetry may not be dead but these latest results have certainly put it into hospital."

(Prof Chris Parkes, spokesperson for the UK Participation in the LHCb experiment)

→Does Not support wimps -CDM-


(In agreement with all dedicated wimp experiments at work from more than 20 years which have not found any wimp's signal) "So far researchers who are racing to find evidence of so called "new physics", ie non-standard models, have run into a series of dead ends".


ANTIMATTER IN SPACE - AMS on board ISS Alpha

Magnet Spectrometer









Positron excess in cosmic rays are not related to DM physics but to astrophysical sources and astrophysical mechanisms and can be explained by them

LHC AMS PLANCK

Three beautiful and big experiments of performant instruments, technology, industry, achievements and successful operation which do not find the main scientific objective emphasized by them (for which they were designed)


• Why No Experimental Detection of the DM particle has been reached so far ?

• Because:

- All experimental searches for DM particles are dedicated to CDM: wimps of m > 1 GeV,
- While the DM particle mass is in the keV scale.
- Moreover, past, present and future reports of signals of such CDM experiments **cannot be due to DM** because of the same reason.
- The inconclusive signals in such experiments should be originated by phenomena of other kinds.
- In addition, such signals contradict each other supporting the idea that they are unrelated to any DM detection.

LUX Large Underground Xenon Detector 30 October 2013

Dark Matter Experiment Has Detected Nothing, Researchers Say Proudly



- They found no sign of WIMPS signals. beyond the expected background noise.
- The experiment did so at far better sensitivities than any such experiment before it.

Sterile Neutrinos ν

- Rhenium and Tritium beta decay (MARE, KATRIN). Theoretical analysis: H J de V, O. Moreno, E. Moya de Guerra, M. Ramón Medrano, N. Sánchez, Nucl. Phys. B866, 177 (2013).
- [Other possibility to detect a sterile ν_s : a precise measure of nucleus recoil in tritium beta decay.]
- Conclusion: the empty slot of right-handed neutrinos in the Standard Model of particle physics can be filled by keV-scale sterile neutrinos describing the DM.

An appealing mass neutrino hierarchy appears:

- ▲ Active neutrino: ~ mili eV
- Light sterile neutrino: ~ eV
- Dark Matter: ~ keV
- ullet Unstable sterile neutrino: \sim MeV....

LHC AMS PLANCK

Three beautiful and big experiments of performant instruments, technology, industry, achievements and successful operation which do not find the main scientific objective emphasized by them (for which they were designed)

Recent News on Cosmological Observables

Before 2013: Hubble constant $H_0=73.8\pm2.4~{
m km\over s}~{1\over
m Mpc}$ from

A G Riess et al. ApJ 730, 119 (2011).

direct observations of Cepheids by HST, $\Omega_m = 0.27 \pm 0.03$

Planck assumed here only three neutrinos and no sterile neutrinos ν_s .

Planck 2013: $H_0 = 67.3 \pm 1.2 \frac{\text{km}}{\text{s}} \frac{1}{\text{Mpc}}$. $\Omega_m = 0.32 \pm 0.02$.

There is today strong evidence for ν_s from short baseline experiments (reactors, MiniBoone, LSND).

Adding one ν_s yields:

 $H_0=70\pm 1.2~{
m km\over s}~{1\over {
m Mpc}}.~\Omega_m=0.30\pm 0.01~{
m for}~m_s=0.4~{
m eV}.$ These values for H_0 and Ω_m are compatible with the direction

astronomical measurements.

M. Wyman et al. arXiv:1307.7715, J. Hamann & J.

Haserkamp, arXiv:1308.3255, R. Battye & A. Moss,

Effective Theory of Inflation (ETI) confirmed by Planck

Quantity	ETI Prediction	Planck 2013
Spectral index $1 - n_s$	order $1/N = 0.02$	0.04
Running $dn_s/dlnk$	order $1/N^2 = 0.0004$	< 0.01
Non-Gaussianity f_{NL}	order $1/N = 0.02$	< 6
	ETI + WMAP+LSS	
tensor/scalar ratio r	r = 0.04-0.05	< 0.11
inflaton potential		
curvature $V''(0)$	V''(0) < 0	V''(0) < 0

ETI + WMAP+LSS means the MCMC analysis combining the ETI with WMAP and LSS data. Such analysis calls for an inflaton potential with negative curvature at horizon exit. The double well potential is favoured (new inflation). D. Boyanovsky, C. Destri, H. J. de Vega, N. G. Sanchez, arXiv:0901.0549, IJMPA 24, 3669-3864 (2009).

Two key observable numbers: associated to the primordial density and primordial gravitons:

$$n_s = 0.9608$$
, r


PREDICTIONS

r < 0.053

r > 0.021

0.021 < r < 0.053

Most probable value: r ~ 0.051



THE PRIMORDIAL COSMIC BANANA

The tensor to scalar ratio r (primordial gravitons) versus the scalar spectral index n_s. The amount of r is always non zero H.J. de Vega, C. Destri, N.G. Sanchez, Annals Phys 326, 578(2011)

The Energy Scale of Inflation


Grand Unification Idea (GUT)

- Penormalization group running of electromagnetic, weak and strong couplings shows that they all meet at $E_{GUT} \simeq 2 \times 10^{16} \; \mathrm{GeV}$
- ullet Neutrino masses are explained by the see-saw mechanism: $m_
 u\sim rac{M_{
 m Fermi}^2}{M_R}$ with $M_R\sim 10^{16}$ GeV.
- Inflation energy scale: $M \simeq 10^{16}$ GeV.

Conclusion: the GUT energy scale appears in at least three independent ways.


Moreover, moduli potentials: $V_{moduli} = M_{\rm SUSY}^4 \ v \left(\frac{\phi}{M_{Pl}} \right)$ ressemble inflation potentials provided $M_{\rm SUSY} \sim 10^{16}$ GeV. First observation of SUSY in nature??

WMAP 5 years data set plus other CMB data




These Acoustic Oscillations are excited by the primordial inflationary power: $P(k) = \Delta k^{n_s-1}$, $n_s =$ spectral index. An explanation for the the quadrupole suppression: the fast-roll stage of inflation. DB, HJdV, NGS, PRD74, 123006 and 123007 (2006).

Fast roll Inflation produces the Observed Quadrupole CMB Suppression



D. Boyanovsky, H. J de Vega and N. G. Sanchez, "CMB quadrupole suppression II: The early fast roll stage"
Phys. Rev. D74, 123006 (2006)

COSMIC HISTORY AND CMB QUADRUPOLE SUPPRESSION



Dark Matter Particles

DM particles decouple due to the universe expansion, their distribution function freezes out at decoupling.

The characteristic length scale is the free streaming scale (or Jeans' scale). For DM particles decoupling UR:

$$r_{Jeans} = 57.2 \, \mathrm{kpc} \, \frac{\mathrm{keV}}{m} \, \left(\frac{100}{g_d} \right)^{\frac{1}{3}}$$
, solving the linear Boltz-V eqs. $g_d = \mathrm{number}$ of UR degrees of freedom at decoupling.

- DM particles can freely propagate over distances of the order of the free streaming scale.
- Therefore, structures at scales smaller or of the order of r_{Jeans} are erased.
- The size of the DM galaxy cores is in the ~ 50 kpc scale $\Rightarrow m$ should be in the keV scale (WDM particles).

For neutrinos $m \sim \text{eV HDM particles}$ $r_{Jeans} \sim 60 \text{ Mpc} \Rightarrow \text{NO GALAXIES FORMED}.$

Dark Matter: from primordial fluctuations to Galaxies

Cold (CDM): small velocity dispersion: small structures form first, bottom-up hierarchical growth formation, too heavy (GeV)

*Hot (HDM): large velocity dispersion: big structures form first, top-down, fragmentation, ruled out, too light (eV)

Warm (WDM): "in between", right mass scale, (keV)


AWDM Concordance Model:

CMB + LSS + SSS Observations

DM is WARM and COLLISIONLESS

- CDM Problems:
- > clumpy halo problem", large number of satellite galaxies "satellite problem", overabundance of small structures
- > \ ρ (r) ~ 1 / r (cusp)
- And other problems.....

Linear primordial power today P(k) vs. k Mpc h




 $\log_{10} P(k)$ vs. $\log_{10}[k \ \mathrm{Mpc} \ h]$ for WIMPS, 1 keV DM particles and 10 eV DM particles. $P(k) = P_0 \ k^{n_s} \ T^2(k)$.


P(k) cutted for 1 keV DM particles on scales $\lesssim 100$ kpc.

Transfer function in the MD era from Gilbert integral eq

WDM vs. CDM linear fluctuations today



Box side = 22.6 Mpc. [C. Destri, private communication].



WARM DARK MATTER REPRODUCE

→OBSERVED GALAXY DENSITIES AND VELOCITY DISPERSIONS

→SOLVES the OVERABUNDANCE ("satellite) PROBLEM

->OBSERVED SURFACE DENSITY VALUES OF DARK MATTER DOMINATED GALAXIES

→OBSERVED GALAXY
CORED DENSITY PROFILES: QUANTUM
MECHANICS

Dwarf galaxies as quantum objects

–de Broglie wavelength of DM particles $\lambda_{dB}=rac{\hbar}{m\,\sigma}$

d = mean distance between particles,

 $\sigma = \mathsf{DM}$ mean velocity

$$d=\left(rac{m}{
ho}
ight)^{rac{1}{3}}$$
 , $Q=
ho/\sigma^3$, $Q=$ phase space density.

ratio:
$$\mathcal{R} = \frac{\lambda_{dB}}{d} = \hbar \left(\frac{Q}{m^4}\right)^{\frac{1}{3}}$$

Observed values:
$$2 \times 10^{-3} < \mathcal{R} \left(\frac{m}{\text{keV}} \right)^{\frac{1}{3}} < 1.4$$

The larger R is for ultracompact dwarfs.

The smaller R is for big spirals.

 \mathcal{R} near unity (or above) means a QUANTUM OBJECT.

Observations alone show that compact dwarf galaxies are quantum objects (for WDM).

The quantum radius r_q for different kinds of DM

DM type	DM particle mass	r_q	
CDM	$1-100~{\sf GeV}$	$1-10^4~{ m meters}$	in practice zero
WDM	1-10~keV	0.1 - 1 pc	compatible with observed cores
HDM	$1-10~\mathrm{eV}$	kpc - Mpc	too big !

RESULTS

All the obtained density profiles are cored.

The Core Sizes are in agreement with the observations

from the compact galaxies where $r_h \sim 20$ pc till the spiral and elliptical galaxies where $r_h \sim 0.2$ - 60 kpc.


The larger and positive is the chemical potential $\nu(0)$, the smaller is the core. The minimal one arises in the degenerate case $\nu(0)$ --> to +infinity (compact dwarf galaxies).

And
The Phase-space Density
The Galaxy halo Masses.

Agreement is found in all the range of galaxies for a DM particle mass m around 2 keV.

Error bars of the observational data are not shown but they are at least about 10-20 %.


vs. halo radius. Galaxy observations vs. Thomas-Fermi



observed $Q = \rho/\sigma^3$ from stars are upper bounds for DM Q

Density and velocity profiles from Thomas-Fermi

Cored density profile and velocity profile obtained from Thomas-Fermi.



THE MINIMAL GALAXY MASS

A minimal galaxy mass and minimal velocity dispersion are found.

This in turn implies a minimal mass m_min =1.91 keV for the WDM particle.

This minimal WDM mass is a universal value, independent of the WDM particle physics model because only relies on the degenerate quantum fermion state, which is universal whatever is the non-degenerate regime.

These results and the observed halo radius and mass of the compact galaxies also provide further indication that the WDM particle mass m is approximately around 2 keV.

More precise data will make this estimation more precise.

Minimal galaxy mass from degenerate WDM

The halo radius, the velocity dispersion and the galaxy mass take their minimum values for degenerate WDM:

$$r_{h \ min} = 24.51 \dots \text{ pc } \left(\frac{m}{\text{keV}}\right)^{\frac{4}{3}} \left[\rho(0) \frac{\text{pc}^{3}}{M_{\odot}}\right]^{\frac{1}{6}}$$
 $M_{min} = 2.939 \dots 10^{5} \ M_{\odot} \left(\frac{\text{keV}}{m}\right)^{4} \sqrt{\rho(0) \frac{\text{pc}^{3}}{M_{\odot}}}$
 $\sigma_{min}(0) = 2.751 \dots \frac{\text{km}}{\text{s}} \left(\frac{\text{keV}}{m}\right)^{\frac{4}{3}} \left[\rho(0) \frac{\text{pc}^{3}}{M_{\odot}}\right]^{\frac{1}{3}}$.

These minimum values correspond to the observations of compact dwarf galaxies.

Lightest known compact dwarf galaxy is Willman I:

$$M_{Willman\ I} = 2.9\ 10^4\ M_{\odot}$$

Imposing $M_{Willman\ I} > M_{min}$ yields the lower bound for the WDM particle mass: m > 1.91 keV.

WARM DARK MATTER REPRODUCE

→OBSERVED GALAXY DENSITIES AND VELOCITY DISPERSIONS

→OBSERVED GALAXY CORED DENSITY PROFILES

->OBSERVED SURFACE DENSITY VALUES OF DARK MATTER DOMINATED GALAXIES

→SOLVES the OVERABUNDANCE ("satellite)
PROBLEM and the CUSPS vs CORES Problem

WDM OVERALL CONCLUSION

• To conclude, we find it is highly remarkable that in the context of warm dark matter, the quantum description provided by this semiclassical framework, (quantum WDM and classical gravitation), is able to reproduce such broad variety of galaxies.

 The resulting galaxy, halo radius, galaxy masses and velocity dispersion are fully consistent with observations for all different types of galaxies. Fermionic WDM treated quantum mechanically, as it must be, is able to reproduce the observed galactic cores and their sizes. In addition, WDM simulations produce the right DM structures in agreement with observations for scales > kpc.

IN PROGRESS

H. J. de Vega, N. G. Sanchez: BLACK HOLES FORMED by WDM and BARYONS

(GALACTIC SUPERMASSIVE, STELLAR)

Galaxy Structure from Classical Cosmological Boltzmann-Vlasov equations:

Generalized Larson equations

And other results.....

keV Sterile Neutrino Warm Dark Matter


Sterile neutrinos can decay into an active-like neutrino and a monochromatic X-ray photon with an energy half the mass of the sterile neutrino. Observing the X-ray photon provides a way to observe sterile neutrinos in DM halos.

WDM keV sterile neutrinos can be copiously produced in the supernovae cores. SN stringently constrain the neutrino mixing angle squared to be 10⁻⁹ for m > 100 keV (in order to avoid excessive energy lost) but for smaller masses the SN bound is not so direct. Within the models worked out till now, mixing angles are essentially unconstrained by SN in the keV mass range.

Sterile neutrinos are produced out of thermal equilibrium and their production can be non-resonant (in the absence of lepton asymmetries) or resonantly enhanced (if lepton asymmetries are present).







How to detect sterile neutrinos?

Sterile neutrinos can be detected in beta decay and in electron capture (EC) when a u_s with mass in the keV sca is produced instead of an active u_e .

Beta decay: the electron spectrum is slightly modified at energies around the mass (\sim keV) of the ν_s .

$$^3H_1 \Longrightarrow {}^3He_2 + e^- + \bar{\nu}_e$$
 , $^{187}Re \Longrightarrow {}^{187}Os + e^- + \bar{\nu}_e$.

The electron energy spectrum is observed.

Electron capture:
$$^{163}Ho + e^{-} \Longrightarrow ^{163}Dy^* + \nu_e$$

The nonradiative de-excitation of the Dy^* is observed and different for u_s in the keV range than for active u_e .

Experiments that may detect sterile neutrinos:

MARE (Milano), KATRIN (Karlsruhe), PTOLEMY (Princeton), ECHo (Heidelberg).

They search the mass of the ordinary neutrino.

END

THANK YOU FOR YOUR ATTENTION

He was a pressured to astrophysics. passion, for his apparimental and channel call to other a sed for the conception and construction of perintermedic, like the bydrogen tobeselection of crophotocease:

Z' a late un precuratre negli a ladi di corregiona, per i cue teneri mental e a perimendali e per aver ideato e replágado esperiatromento quelli il tubo a lidrogeno e il migrofolometro



Dispisi. Challonge (1895-1977), stirromonar of the Observatoire de Parin and one of the foundati of the Intrinsi d'Antrophytique de Paria, dedicated his life at evo postions: the adestilla repeatable de-

Davidi. Citalongo (1895-1977), astronomo dell'Observatoire de Part, e uno dei fondatori dell'Issira. di Ja traficias di Parigi, ka dediasto la vita a due passioni. La riceras scientifica e la montagna.

Els was a piocese lo des escencis al Eligio Aldiande O bouvaionise, si Pic do Elife and al Bongfranjoch

El photo wegli avui. 30 un pionere delle ricarche pretto gli Ott er arori di Airo Mantagno, où lezenvolente del . He du Mild e où a zienjeue di nierrez di Angfreujech.

He gave his cause to an explayable tentangues of the categophotocastic and the operatographs Chategop and to a periodo the Alpo -the Peak Chalcoge -.

Na data il suo some a strumenti di astrofisias -il microfisometro, e los pritragrafi Chalouge, e a : una cima delle Alpi -è picas Challong-...

Today the International School of Astrophysics Daniel Chalcogs, discond by Professor Flores. Sancher -Observatoire de Paris-La camedation bira.

Oggi è indibilata a suo vorre la livier rational School of la traphysia: David Challouge, diretto dall Professor Horma Sancheg -Observatoire de Paris-




nica Chilaga Lautani. 1934, 1935



Temperaturi des primes régimitaises des distribualde adole Alberta 1988)





"... The effects required for studying the stronge phenomena at the extreme border of the Earth and the great services provided from rile high mesuurole eitser sories..."

". . Gil a forg i ridicio di per la acudia degli arrani favorenzi dila a vengava ai auglisi della terra e gli limmetali autoggi die si persone estimator con in es servações i la cita in estagac..."

David Challege, La Maurager et l'alpiet ver 19245, 1988







The quality sph Chin egosi Feeglessjoch



de l'anglement with the 25 and descripe

Donald Challenge of Feeglinshale

Dani di Graf enge hauparta la spattagrafa nationalise à d'Onnervations 😋 bien (Desgles) protestituente e como protesti di Tampane di 100 m.

> En queliques heures de mardie sur les senders des Aligno, pome i ino unigno el ino nechero, je vio pilos, le requelle plus d'insure plans, plus de pour entre que en aix maia de la via . .

Dawiel Challenge

In fine hours of welling on the cips treats, between the sum and the racks, Leother more emotions, mare memories than in six months of normal life...

De papite are de comunica sui seuriere delle Alpi, irale se de le recer, recesige plu se exequed, plu ricardi cile is sei mesi di vis...

At that time -first half of 1900- also the salentific research was more interes and more effective to the High Manuscia. Observatories.

A quel ten pe-prima metà del 1900audio la ricerca salestifica era più interes and officere in disc manages.







Fig. 12. — Maquette de la surface Σ .

Daniel Chalonge et Jean Prouvé : la rencontre de l'œuvre de deux pionniers après leur temps



Jean Prouvé et l'Observatoire de Paris Quartier de l'Observatoire.

Depuis « les maisons industrialisées de Meudon au Bat méridien de l'instrument des Passages À » l'Observatoire de Paris






Le bat Prouvé situé juste dans le terrain de transition entre l'Observatoire de Paris et l'IAP Daniel Chalonge a fait précisément le chemin, la transition, entre l'Observatoire de Paris et l'IAP, puisque il à été un des fondateurs de l'IAP, passage de l'astronomie à l'astrophysique

Le fait que aujourd'hui les archives et instruments Chalonge sont collectés dans le bat Prouvé (qui est un hasard des circunstances) rend hommage à la mémoire « in situ » de Daniel Chalonge, et de Jean Prouvé, car il rend la vocation astronomique/astrophysique à ce bâtiment , à cette partie du terrain.et à posteriori de Jean Prouvé.

École Internationale Daniel Chalonge





22 Years of Activity Calling for Understanding

SCIENCE WITH GREAT INTELLECTUAL ENDEAVOUR AND A HUMAN FACE LA SCIENCE QUI DONNE ENVIE : UNE GRANDE AVENTURE SCIENTIFIQUE ET HUMAINE

PROGRAMME 2013

15 MARCH 2013 : "Présentation du Programme 2013 et des Dernières Nouvelles Scientifiques de l'Univers » Bâtiment Perrault, Observatoire de Paris

4-7 APRIL 2013: "Latest News from the Universe, Dark Matter Galaxies and Particle Physics" Palazzo de l'Università & Palazzo Graneri, Piamonte Région, Turin, Italy

16 MAY 2013: Spring Open Session of Scientific Culture 2013 Session Ouverte de Printemps de Culture Scientifique 2013 : "L'Homme et l'Univers" Bâtiment Perrault, Observatoire de Paris, Paris

30 MAY 2013: Rencontre de Culture Scientifique "Voyage à travers l'Univers: De ses Origines à nos Jours" Cité Internationale Universitaire de Paris, Paris

4-7 JUNE 2013 : Chalonge Meudon Workshop 2013 "Warm Dark Matter Galaxies in Agreement with Observations: Formation, Evolution and Supermassive Black Holes" Observatoire de Paris, Château de Meudon-CIAS, Meudon

23-26 JULY 2013: The 17th Paris Cosmology Colloquium Chalonge 2013: "The New Standard Model of the Universe: AWDM - Warm Dark Matter: "Theory and Observations" Bâtiment Perrault, Observatoire de Paris, Paris

26 JULY 2013: Summer Open Session of Scientific Culture 2013 / Session Ouverte d'Eté de Culture Scientifique 2013 : A Surprise Session

AUTOMME 2013: Cycle Les grandes questions posées aujourd'hui à la Science: 1ère Question : Où va la Science ? Cité Internationale Universitaire de Paris, Paris

And Other Events...

N. G. SANCHEZ * H. J. DE VEGA * M. C. FALVELLA * A. ZANINI * M. RAMON MEDRANO * A. PERISSA and other colleagues http://chalonge.obspm.fr













La Science qui donne envie. Une grande aventure scientifique et humaine



14 MARCH 2014 : Opening Session 2014. Session ouverte de Culture Scientifique "Présentation du Programme 2014 et des Dernières Nouvelles Scientifiques de l'Univers" Bâtiment Perrault, Observatoire de Paris

22 MAY 2014: Spring Open Session of Scientific Culture 2014
Session Ouverte de Printemps de Culture Scientifique Interdisciplinaire 2014:
"L'Homme et l'Univers" Bâtiment Perrault, Observatoire de Paris, Paris.

4-7 JUNE 2014 : Chalonge Meudon Workshop 2014 "From large to small scale structures in agreement with observations: CMB, WDM, galaxies, black holes, neutrinos and sterile neutrinos" Observatoire de Paris, Château de Meudon-CIAS, Meudon

22-25 JULY 2014: The 18th Paris Cosmology Colloquium Chalonge 2014: "Latest News from the Universe: LambdaWDM, CMB, Dark Matter, Dark Energy, Neutrinos and Sterile Neutrinos" Bâtiment Perrault, Obs.de Paris

25 JULY 2014 : Summer Open Session of Scientific Culture 2014 Session Ouverte d'Eté de Cultur e Scientifique 2014 : A Surprise Session

AUTOMME 2014: Cycle Les grandes questions posées aujourd'hui à la Science: Où va la Science ? Cité Internationale Universitaire de Paris, Paris

17-18 OCTOBRE 2014: Chalonge Turin Session 2014 "Latest News from the Universe, Dark Matter Galaxies and Particle Physics" Palazzo de l'Università & Accademia delle Scienze, Piamonte Région, Turin, Italy.

27-28 NOVEMBER 2014 : Concluding Session 2014 & Avant-Première 2015
And Other Events... http://chalonge.obspm.fr

Welcome to the Chalonge School: A Laboratory of Ideas. Research. Training. Scientific Culture.

.....

A beacon pioneering and developping research, projects and training.

The programme offers unvaluable international current research view at the forefront of astrophysics and cosmology, international contacts at the highest level and a careful interdisciplinarity, with both Theory and Observations.

The programme is open to researchers, post-docs and advanced students of the different disciplines in the field, both theorists, experimentalists, observers. Advanced students, post-docs, young researchers are encouradged to participate. The programme includes scientific culture events with the latest results and exhibitions.

The Chalonge School Team

Organizers:

Norma G. SANCHEZ, Héctor J. DE VEGA, Maria C. FALVELLA, Alba ZANINI, Marina RAMON MEDRANO, Annalisa PERISSA and other colleagues

Technical Support:

Djilali ZIDANI, François SEVRE, Nicole LETOURNEUR, Jean-Pierre MICHEL, Sylvain CNUDDE, Emmanuel VERGNAUD, Jèrome BERTHIER, and other colleagues....

The Chalonge School Medal

The Chalonge Medal is coined exclusively for the Chalonge School by the Hôtel de la Monnaie de Paris (the French Mint). Only ten Chalonge medals have been awarded in the 22 year school history.

Awarded Daniel Chalonge Medals:

The Chalonge School Team

Organizers:

Norma G. SANCHEZ, Héctor J. DE VEGA, Maria C. FALVELLA, Alba ZANINI, Marina RAMON MEDRANO, Annalisa PERISSA and other colleagues

Technical Support:

Djilali ZIDANI, François SEVRE, Nicole LETOURNEUR, Jean-Pierre MICHEL, Sylvain CNUDDE, Emmanuel VERGNAUD, Jèrome BERTHIER, and other colleagues....

The Chalonge School Medal

The Chalonge Medal is coined exclusively for the Chalonge School by the Hôtel de la Monnaie de Paris (the French Mint). Only ten Chalonge medals have been awarded in the 22 year school history.

Awarded Daniel Chalonge Medals:

Subramanyan CHANDRASEKHAR, Nobel prize of physics. Bruno PONTECORVO.

George SMOOT, Nobel prize of physics.

Carlos FRENK.

Anthony LASENBY.

Bernard SADOULET, Fellow of the USA Academy of Arts And Sciences.

Peter BIERMANN.

John MATHER, Nobel prize of physics.

Brian SCHMIDT, Nobel prize of Physics.

Gérad GILMORE, Fellow of the UK Royal Society.

.....

School Courses, Lectures and Lecturers, Album of Pictures: http://chalonge.obspm.fr